

# Electromagnetic Transition Rate Studies with the TIGRESS Integrated Plunger

P. Voss<sup>1</sup>, K. Starosta<sup>1</sup>, C. Andreoiu<sup>1</sup>, R. Ashley<sup>1</sup>, R. Austin<sup>2</sup>, G. Ball<sup>3</sup>, A. Chester<sup>1</sup>, A. B. Garnsworthy<sup>3</sup>, P. Garrett<sup>4</sup>, G. Hackman<sup>3</sup>, R. Henderson<sup>3</sup>, R. Krücken<sup>3</sup>, C. Svensson<sup>4</sup>, and The TIGRESS Collaboration

<sup>1</sup>Simon Fraser University, Department of Chemistry, Burnaby, BC, Canada V5A 1S6

<sup>2</sup>St. Mary's University, Department of Astronomy and Physics, Halifax, NS, Canada B3H 3C3

<sup>3</sup>TRIUMF, Vancouver, BC, Canada, V6T 2A3


<sup>4</sup>University of Guelph, Department of Physics, Guelph, ON, Canada N1G 2W1

SFU

## Introduction

- Precise transition rate measurements are fundamental probes of nuclear structure and permit stringent tests for theoretical models important to our understanding of the nucleus.
- The TIGRESS Integrated Plunger (TIP) delivers a new experimental program for transition rate studies via Doppler-shift lifetime and sub-barrier Coulomb excitation measurements using accelerated radioactive beams from the ISAC-II facility at TRIUMF.
- Experimental and developmental work on TIP and its extensive suite of ancillary charged-particle detector systems has been focused on probing nuclear structure along the  $N=Z$  line.
  - Impact of increasing neutron deficiency on shape evolution of medium mass nuclei.
  - Quality of shell model descriptions of the properties of low-lying levels in light nuclei.




## The TIGRESS Integrated Plunger



- Plunger gamma-ray spectra are collected with TIGRESS for recoil distance method lifetime measurements.
- TIGRESS is an array of HPGe segmented clover detectors with BGO suppressors and fast digital electronics.
- Future TIP lifetime studies of weak and exotic reaction channels will benefit from new spectroscopy tools: EMMA (ElectroMagnetic Mass Analyser) and DESCANT (Deuterated Scintillator Array for Neutron Tagging).

## Transition Rate Discrepancies in <sup>36</sup>Ar


Doppler-shift attenuation method (DSAM) lifetime and sub-barrier Coulomb excitation data have been collected with the TIP target wheel and ancillary silicon detector systems to help resolve recent transition rate discrepancies in <sup>36</sup>Ar.



DSAM schematic for stopper of thickness  $d = v \cdot t_{\text{stopping}}$

## Nuclear Shape Evolution and <sup>68</sup>Se

- Along the  $N=Z$  line, protons and neutrons occupy the same shell model orbitals. The large wave function overlap leads to an amplification of their interactions, which drives a steady deformation of nuclear shapes between <sup>56</sup>Ni and <sup>100</sup>Sn.
- $B(E2)$  transition strength model calculations vary by an order of magnitude. Data (shown below from Ref. [4]) from intermediate energy Coulomb excitation indicate the shape evolution from <sup>64</sup>Ge to <sup>68</sup>Se is suppressed.
- TIP will provide a direct, model-independent, and complimentary transition rate measurement following fusion-evaporation reactions via Doppler-shift lifetime measurements.





- The <sup>40</sup>Ca(<sup>36</sup>Ar, 2α)<sup>68</sup>Se reaction channel will be selected with evaporated light-particle detection using the TIP CsI(Tl) ball, which will offer nearly 3π coverage.
- Integration of the CsI array into the TIGRESS DAQ will permit pulse-shape analysis of the collected waveforms to enable reaction channel selectivity.



Various stages of development of the TIP CsI ball.

## Present Analysis and Results from TIP



- [Left] <sup>36</sup>Ar beam energies of 115, 130, 145, and 160 MeV were delivered onto a <sup>40</sup>Ca target and the fusion-evaporation reaction channels were identified by gamma-ray energies. A qualitative analysis of the excitation function was performed and compared to PACE4 calculations to determine the optimal <sup>68</sup>Se production beam energy.
- [Middle] Prototype CsI(Tl) crystals coupled to silicon PIN diodes were used for evaporated charged-particle identification via pulse-shape analysis of the excitation function data.
- [Right] Gamma-ray decay energy spectra at forward (45°) and backward (135°) TIGRESS angles following the Coulomb excitation of <sup>36</sup>Ar on a gold-backed carbon target. Gamma-rays were collected in coincidence with recoiling carbon nuclei.

## Summary

- Electromagnetic transition rate measurements with accelerated radioactive beams play an important role in our understanding of nuclear structure.
- The TIGRESS Integrated Plunger and associated ancillary detector systems offer a high degree of flexibility for precise Doppler-shift lifetime and sub-barrier Coulomb excitation studies along the  $N=Z$  line.
- Experimental and analytical efforts towards resolving discrepancies in the self-conjugate <sup>36</sup>Ar system are underway.
- Work continues to fully develop TIP and the CsI array for transition rate studies of exotic nuclei far from stability.

## Acknowledgements

We thank the TIGRESS collaboration and the SFU Science Machine and Electronics Shops for their indispensable assistance.

- [1] K.-H. Speidel et al., Phys. Lett. B **632**, 207 (2006).
- [2] B. V. Pritychenko et al., Phys. Lett. B **461**, 322 (1999).
- [3] P. D. Cottle et al., Phys. Rev. C **60**, 031301(R) (1999).
- [4] A. Obertelli et al. Phys. Rev. C **80**, 031304(R) (2009).

